

On the Companions of Argon

Author(s): William Ramsay and Morris W. Travers

Source: *Proceedings of the Royal Society of London*, Vol. 63 (1898), pp. 437-440

Published by: The Royal Society

Stable URL: <http://www.jstor.org/stable/116011>

Accessed: 14/08/2009 06:02

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=rsl>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



The Royal Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the Royal Society of London*.

investigated by means of radiometer mill wheels, recently described in a paper to the Physical Society.

In any case, it seems clear that in the tubes observed and photographed with the pin-hole camera, the Röntgen rays given off by certain portions of the fluorescent glass are not originated by the impact of an ordinary cathode stream, but apparently by the impact of positively charged streams proceeding from the anti-cathode.

The writer is greatly indebted to Mr. J. C. M. Stanton and Mr. H. Tyson Wolff, for the construction of the apparatus described, as also for valuable assistance in the carrying out of the experiments.

“On the Companions of Argon.” By WILLIAM RAMSAY, F.R.S., and MORRIS W. TRAVERS. Received June 13,—Read June 16, 1898.

For many months past we have been engaged in preparing a large quantity of argon from atmospheric air by absorbing the oxygen with red-hot copper, and the nitrogen with magnesium. The amount we have at our disposal is some 18 litres. It will be remembered that one of us, in conjunction with Dr. Norman Collie, attempted to separate argon into light and heavy portions by means of diffusion, and, although there was a slight difference* in density between the light and the heavy portions, yet we thought the difference too slight to warrant the conclusion that argon is a mixture. But our experience with helium taught us that it is a matter of the greatest difficulty to separate a very small portion of a heavy gas from a large admixture of a light gas; and it therefore appeared advisable to re-investigate argon, with the view of ascertaining whether it is indeed complex.

In the meantime, Dr. Hampson had placed at our disposal his resources for preparing large quantities of liquid air, and it was a simple matter to liquify the argon which we had obtained by causing the liquid air to boil under reduced pressure. By means of a two-way stopcock the argon was allowed to enter a small bulb, cooled by liquid air, after passing through purifying reagents. The two-way stopcock was connected with mercury gas-holders, as well as with a Töpler pump, by means of which any part of the apparatus could be thoroughly exhausted. The argon separated as a liquid, but at the same time a considerable quantity of solid was observed to separate partially round the sides of the tube, and partially below the

* Density of lighter portion, 19·93; of heavier portion, 20·01, ‘Roy. Soc. Proc.’, vol. 60, p. 206.

surface of the liquid. After about 13 or 14 litres of the argon had been condensed, the stopcock was closed, and the temperature was kept low for some minutes in order to establish a condition of equilibrium between the liquid and vapour. In the meantime, the connecting tubes were exhausted and two fractions of gas were taken off by lowering the mercury reservoirs, each fraction consisting of about 50 or 60 cubic cm. These fractions should contain the light gas. In a previous experiment of the same kind, a small fraction of the light gas had been separated, and was found to have the density 17·2. The pressure of the air was now allowed to rise, and the argon distilled away into a separate gas-holder. The white solid which had condensed in the upper portion of the bulb did not appear to evaporate quickly, and that portion which had separated in the liquid did not perceptibly diminish in amount. Towards the end, when almost all the air had boiled away, the last portions of the liquid evaporated slowly, and when the remaining liquid was only sufficient to cover the solid, the bulb was placed in connection with the Töpler pump, and the exhaustion continued until the liquid had entirely disappeared. Only the solid now remained, and the pressure of the gas in the apparatus was only a few millimetres. The bulb was now placed in connection with mercury gas-holders, and the reservoirs were lowered. The solid volatilised very slowly, and was collected in two fractions, each of about 70 or 80 cubic cm. Before the second fraction had been taken off, the air had entirely boiled away, and the jacketing tube had been removed. After about a minute, on wiping off the coating of snow with the finger, the solid was seen to melt, and volatilise into the gas-holder.

The first fraction of gas was mixed with oxygen, and sparked over soda. After removal of the oxygen with phosphorus it was introduced into a vacuum-tube, and the spectrum examined. It was characterised by a number of bright red lines, among which one was particularly brilliant, and a brilliant yellow line, while the green and the blue lines were numerous, but comparatively inconspicuous. The wave-length of the yellow line, measured by Mr. Baly, was 5849·6, with a second-order grating spectrum. It is, therefore, not identical with sodium, helium, or krypton, all of which equal it in intensity. The wave-lengths of these lines are as follows:—

Na (D ₁)	5895·0
Na (D ₂)	5889·0
He (D ₃)	5875·9
Kr (D ₄)	5866·5
Ne (D ₅)	5849·6

The density of this gas, which we propose to name "neon"

(new), was next determined. A bulb of 32·35 cubic cm. capacity was filled with this sample of neon at 612·4 mm. pressure, and at a temperature of 19·92° it weighed 0·03184 gram.

Density of neon 14·67.

This number approaches to what we had hoped to obtain. In order to bring neon into its position in the periodic table, a density of 10 or 11 is required. Assuming the density of argon to be 20, and that of pure neon 10, the sample contains 53·3 per cent. of the new gas. If the density of neon be taken as 11, there is 59·2 per cent. present in the sample. The fact that the density has decreased from 17·2 to 14·7 shows that there is a considerable likelihood that the gas can be further purified by fractionation.*

That this gas is a new one is sufficiently proved, not merely by the novelty of its spectrum and by its low density, but also by its behaviour in a vacuum-tube. Unlike helium, argon, and krypton, it is rapidly absorbed by the red-hot aluminium electrodes of a vacuum-tube, and the appearance of the tube changes, as pressure falls, from fiery red to a most brilliant orange, which is seen in no other gas.

We now come to the gas obtained by the volatilisation of the white solid which remained after the liquid argon had boiled away.

When introduced into a vacuum-tube it showed a very complex spectrum, totally differing from that of argon, while resembling it in general character. With low dispersion it appeared to be a banded spectrum, but with a grating, single bright lines appear, about equidistant throughout the spectrum, the intermediate space being filled with many dim, yet well-defined lines. Mr. Baly has measured the bright lines, with the following results. The nearest argon lines, as measured by Mr. Crookes, are placed in brackets:—

Reds very feeble, not measured.		
First green band, first bright line	5632·5	(5651 : 5619)
" second " 	5583·0	(5619 : 5567)
" third " 	5537·0	(5557 : 5320)
Second green band, first bright line . . .	5163·0	(5165)
" second " 	5126·5	(5165 : 5065) brilliant.
First blue band, first bright line	4733·5	(4879)
" second " 	4711·5	(4701)
Second blue band, first bright line.	4604·5	(4629 : 4594)
Third blue band (first order)	4314·0	(4333 : 4300)
Fourth blue band (second order)	4213·5	(4251 : 4201)
Fifth blue band (first order), about	3878	(3904 : 3835)

The red pair of argon lines were faintly visible in the spectrum. The density of this gas was determined with the following

* June 16th. After fractionation of the neon, the density of the lightest sample had decreased to 13·7.

results :—A globe of 32·35 c.c. capacity, filled at a pressure of 765·0 mm., and at the temperature 17·43°, weighed 0·05442 gram. The density is therefore 19·87. A second determination, made after sparking, gave no different result. This density does not sensibly differ from that of argon.

Thinking that the gas might possibly prove to be diatomic, we proceeded to determine the ratio of specific heats :—

Wave-length of sound in air	34·18
" " gas	31·68
Ratio for air	1·408
" gas	1·660

The gas is therefore monatomic.

Inasmuch as this gas differs very markedly from argon in its spectrum, and in its behaviour at low temperatures, it must be regarded as a distinct elementary substance, and we therefore propose for it the name "metargon." It would appear to hold the position towards argon that nickel does to cobalt, having approximately the same atomic weight, yet different properties.

It must have been observed that krypton does not appear during the investigation of the higher-boiling fraction of argon. This is probably due to two causes. In the first place, in order to prepare it, the manipulation of a volume of air of no less than 60,000 times the volume of the impure sample which we obtained was required; and in the second place, while metargon is a solid at the temperature of boiling air, krypton is probably a liquid, and more volatile at that temperature. It may also be noted that the air from which krypton has been obtained had been filtered, and so freed from metargon. A full account of the spectra of those gases will be published in due course by Mr. E. C. C. Baly.

"Summary of the principal Results obtained in a Study of the Development of the Tuatara (*Sphenodon punctatum*)."
By ARTHUR DENDY, D.Sc., Professor of Biology in the Canterbury College, University of New Zealand. Communicated by Professor G. B. HOWES, F.R.S. Received June 15,—
Read June 16, 1898.

Thanks to the most generous and freely rendered services of Mr. P. Henaghan, Principal Keeper of the Lighthouse on Stephen's Island in Cook Straits, I have lately obtained a very perfect series of Tuatara embryos, ranging in age from just before the appearance of the blastopore to about the time of hatching. I have classified these embryos in sixteen stages, and propose shortly to publish a general